

CONES AND CONVEX BODIES WITH MODULAR FACE LATTICES.

DANIEL LABARDINI-FRAGOSO, MAX NEUMANN-COTO, AND MARTHA TAKANE

Dedicated to Claus M. Ringel on the occasion of his 60th birthday.

ABSTRACT. If a convex body C has modular and irreducible face lattice (and is not strictly convex), there is a face-preserving homeomorphism from C to a section of a cone of hermitian matrices over $\mathbb{R}$, $\mathbb{C}$, or $\mathbb{H}$, or C has dimension 8, 14 or 26.

1. INTRODUCTION.

Let C be a convex body in $\mathbb{R}^n$. A subset F of C is a *face* of C if every open interval in C that contains a point of F is contained in F . An *extreme point* is a 1-point face. If S is any subset of C , the *face generated by S* is the minimal face of C containing S . The set $\mathcal{F}(C)$ of all faces of C ordered by inclusion is a lattice, where $F \wedge G$ is the intersection of F and G , and $F \vee G$ is the face generated by $F \cup G$. The lattice $\mathcal{L}(C)$ is always *algebraic* (the chains of faces are finite), *atomic* (faces are generated by extreme points) and *complemented* (for every face F there exists a face G such that $F \wedge G = \emptyset$ and $F \vee G = C$). We want to consider convex bodies for which $\mathcal{F}(C)$ is *modular*, i.e. $F \vee (G \wedge H) = (F \vee G) \wedge H$ whenever $F \leq H$. Modularity is a ‘weak distributivity’ property satisfied by the lattice of normal subgroups of a group and by the lattice of subspaces of a vector space. For algebraic, atomic lattices, modularity is equivalent to the existence of a rank function such that $rk(F) + rk(G) = rk(F \vee G) + rk(F \wedge G)$ for all F and G [4]. Strictly convex bodies and simplices clearly have modular face lattices. No other polytopes have this property [3], but there are beautiful examples of non-polytopal convex bodies in which every pair of extreme points is contained in a proper face and every pair of faces with more than one point meet.

If $\mathcal{L}_1$ and $\mathcal{L}_2$ are lattices, their *direct product* is given by $(\mathcal{L}_1 \times \mathcal{L}_2, \leq)$, where $(a, b) \leq (c, d)$ if and only if $a \leq c$ and $b \leq d$. It follows that the direct product of two lattices is modular if and only if the factors are modular. A lattice is called *irreducible* if it is not isomorphic to a direct product of two nontrivial lattices.

If $C_1 \subset \mathbb{R}^m$ and $C_2 \subset \mathbb{R}^n$ are convex bodies, define $C_1 * C_2 \subset \mathbb{R}^{m+n+1}$ as the convex hull of a copy of C_1 and a copy of C_2 placed in general position in the sense that their linear spans are disjoint and have no common directions. So $C_1 * C_2$ is well defined up to a linear transformation: it is the convex join of C_1 and C_2 of largest dimension. For example $C * \{pt\}$ is a pyramid with base C . Let’s say that a convex body C is **-decomposable* if $C = C_1 * C_2$ for two convex bodies C_1 and C_2 .

1991 *Mathematics Subject Classification.* 52A20, 06C05, 51A05, 15A48.

Key words and phrases. convex, face lattice, modular, hermitian matrix, projective space
Research partially supported by PAPIIT-UNAM grant IN103508 and a fellowship from PASPA.

The natural correspondence (up to linear transformation) between convex bodies in $\mathbb{R}^n$ and closed cones in $\mathbb{R}^{n+1}$ gives an isomorphism of face lattices in which $C_1 * C_2$ corresponds to the direct product of the cones, so the results of this paper apply to cones. This project started with the undergraduate thesis of D. Labardini-Fragoso [9], who showed that in dimension less than 6 any cone with modular face lattice is strictly convex or is decomposable (this was conjectured by Barker in [3]).

Lemma 1. *A convex body C is $*$ -decomposable if and only if its lattice of faces $\mathcal{L}(C)$ is reducible.*

Proof. Let $C = C_1 * C_2$. Observe that each point p of $C_1 * C_2$ with $p \notin C_i$, lies in a unique segment joining a point p_1 of C_1 and a point p_2 of C_2 . For, if a point lies in two segments $p_1 p_2$ and $p'_1 p'_2$ then the lines $p_1 p'_1$ and $p_2 p'_2$ are parallel or they intersect, contradicting the assumptions on the spans of C_1 and C_2 . Moreover, if a point p moves along a straight line in $C_1 * C_2$ then the corresponding points p_1 and p_2 move along straight lines in C_1 and C_2 : If p and q are points in C and $x \in pq$ then $x = tp + (1-t)q = t\lambda p_1 + t(1-\lambda)p_2 + (1-t)\mu q_1 + (1-t)(1-\mu)q_2$ which can be rewritten as a linear combination of a point in $p_1 q_1$ and a point in $p_2 q_2$ with coefficients adding up to 1 so $x_1 \in p_1 q_1$ and $x_2 \in p_2 q_2$. Now if C'_i is a face of C_i then $C'_1 * C'_2$ is a face of $C_1 * C_2$. For, if $x \in C'_1 * C'_2$ and $x = \lambda p + (1-\lambda)q$ with $p, q \in C'_1 * C'_2$, then x_1 lies in $p_1 q_1$ and x_2 lies in $p_2 q_2$ so as C'_i is a face of C_i , p_i and q_i lie in C'_i so p and q lie in $C'_1 * C'_2$. Conversely, if C' is a face of $C_1 * C_2$ and $p \in C'$ then p_1 and p_2 lie in C' so $C' = (C' \cap C_1) * (C' \cap C_2)$. It remains to show that $C' \cap C_i$ is a face of C_i . If $x \in C' \cap C_1$ and $x = \lambda p + (1-\lambda)q$ with $p, q \in C_1 * C_2$ then as C' , and $C_1 = C_1 * \emptyset$ are faces of $C_1 * C_2$, p and q lie in C' and also in C_1 , so $C' \cap C_1$ is a face of C_1 . $C' \cap C_2$ is a face of C_2 . So $\mathcal{L}(C_1 * C_2) \simeq \mathcal{L}(C_1) \times \mathcal{L}(C_2)$.

If $\mathcal{L}(C) \approx \mathcal{L}_1 * \mathcal{L}_2$ then $\mathcal{L}_1$ and $\mathcal{L}_2$ are isomorphic to sublattices of $\mathcal{L}(C)$, so $\mathcal{L}_i \approx \mathcal{L}(C_i)$ for two faces of C with $C_1 \wedge C_2 = \emptyset$ and $C_1 \vee C_2 = C$. To show that $C = C_1 * C_2$ we need to prove that $\text{span}(C_1)$ and $\text{span}(C_2)$ are disjoint and have no directions in common. Suppose that $x \in \text{span}(C_1) \cap \text{span}(C_2)$. Take $x_i \in \text{Int}(C_i)$ then the line through x and x_i meets ∂C_i at two points a_i and b_i . As a_2 lies in a proper subface C'_2 of C_2 , the face generated by C_1 and a_2 lies in $C_1 \vee C'_2$ which is a proper subface of $C_1 \vee C_2$. But the points a_1, b_1, a_2, b_2 determine a plane quadrilateral whose side $a_i b_i$ lies in the interior of C_i so its diagonals intersect at an interior point c of $C_1 \vee C_2$ so the face generated by C_1 and a_2 (which contains c) must be $C_1 \vee C_2$, a contradiction. Now suppose that $\text{span}(C_1)$ and $\text{span}(C_2)$ have a common direction v . Take $x_i \in \text{Int}(C_i)$ then the line through x_i in the direction v meets ∂C_i at two points a_i and b_i . As before a_1, b_1, a_2, b_2 determine a plane quadrilateral whose diagonals intersect at an interior point c of $C_1 \vee C_2$, but c lies in the face generated by C_1 and a_2 which is a proper face of $C_1 \vee C_2$. $\square$

Recall that a *projective space* consists of a set P (the points) and a set L (the lines) so that (a) Each pair of points is contained in a unique line, (b) If a, b, c, d are distinct points and the lines ab and cd intersect, then the lines ac and bd intersect (c) Each line contains at least 3 points and there are at least 2 lines (d) Every chain of subspaces (also called *flats*) has finite length. The maximum length of a chain starting with a point is the *projective dimension* of the space.

The flats of a projective space form an algebraic, atomic, irreducible, modular lattice. Conversely, any lattice with these properties is the lattice of flats of a

projective space, whose points are atoms and whose lines are joins of two atoms [6]. It is a classic result of Hilbert [8] that a projective space in which Desargues theorem holds is isomorphic to the projective space $\mathbb{A}\mathbb{P}^n$ determined by the linear subspaces of $\mathbb{A}^{n+1}$, for some division ring $\mathbb{A}$, and that $\mathbb{A}\mathbb{P}^n$ and $\mathbb{B}\mathbb{P}^m$ are isomorphic if and only if $\mathbb{A}$ and $\mathbb{B}$ are isomorphic and $m = n$. All projective spaces of dimension larger than 2 are desarguesian, but there are many non-desarguesian projective planes.

Examples of convex bodies whose face lattices determine the projective spaces $\mathbb{R}\mathbb{P}^n$, $\mathbb{C}\mathbb{P}^n$ and $\mathbb{H}\mathbb{P}^n$, and the octonionic projective plane arise as sections of some classical cones.

Example 1. Let $\mathbb{F} \in \{\mathbb{R}, \mathbb{C}, \mathbb{H}\}$, let $H_n(\mathbb{F})$ be the set of Hermitian (self-adjoint) $n \times n$ matrices with coefficients in $\mathbb{F}$, and let $C_n(\mathbb{F})$ be the subset of positive-semidefinite matrices (A is positive-semidefinite if $\overline{v}Av^T \geq 0$, for all $v \in \mathbb{F}^n$). Then $C_n(\mathbb{F})$ is a real cone whose face lattice is isomorphic to the lattice of subspaces of $\mathbb{F}^n$.

To see this, let $A, B \in C_n(\mathbb{K})$, and let $\varphi(B)$ denote the face generated by B . Then $A \in \varphi(B)$ if and only if $\ker A \supseteq \ker B$. For, $A \in \varphi(B) \Leftrightarrow \exists \lambda > 0$ such that $B - \lambda A \in C_n(\mathbb{K}) \Leftrightarrow \exists \lambda > 0$ such that $\overline{w}Bw^T \geq \lambda \overline{w}Aw^T \geq 0$ for all $w \in \mathbb{F}^n \Leftrightarrow \overline{w}Bw^T = 0$ implies $\overline{w}Aw^T = 0$ for all $w \in \mathbb{F}^n \Leftrightarrow \ker A \supseteq \ker B$ (since for $A \in C_n(\mathbb{K})$, $\overline{w}Aw^T = 0$ if and only if $Aw^T = 0$). Therefore $\varphi(A) \rightarrow (\ker A)^\perp$ defines a bijection ν from the set of faces of $C_n(\mathbb{K})$ to the set of linear subspaces of $\mathbb{K}^n$. To prove that ν is an isomorphism of lattices observe that $\nu(\varphi(A) \vee \varphi(B)) = \nu(\varphi(A + B)) = (\ker(A + B))^\perp = (\ker A \cap \ker B)^\perp = (\ker A)^\perp \cup (\ker B)^\perp$, and on the other hand, if $\varphi(A) \wedge \varphi(B)$ is a non-empty face, then it is generated by a matrix C with $\ker C = \text{span}(\ker A \cup \ker B)$, so $\nu(\varphi(A) \wedge \varphi(B)) = (\ker C)^\perp = (\ker A)^\perp \cap (\ker B)^\perp$.

Example 2. Let $H_3(\mathbb{O})$ be the set of Hermitian 3×3 matrices over $\mathbb{O}$. Then the subset $C_3(\mathbb{O})$ of all sums of squares of elements in $H_3(\mathbb{O})$ is a real cone whose face lattice determines an octonionic projective plane.

This can be shown using the nontrivial fact that each matrix in $H_3(\mathbb{O})$ is diagonalizable by an automorphism of $H_3(\mathbb{O})$ that leaves the trace invariant [2], so:

(a) A matrix A in $H_3(\mathbb{O})$ lies in $C_3(\mathbb{O})$ if and only if it can be diagonalized to a matrix A' with non-negative entries, because if A lies in $C_3(\mathbb{O})$ then A' is a sum of squares of matrices in $H_3(\mathbb{O})$, which have non-negative diagonal entries.

(b) All the idempotent matrices in $H_3(\mathbb{O})$ lie in $C_3(\mathbb{O})$ as they are squares ($A = A^2$). The idempotent matrices of trace 1 correspond to the extreme rays of $C_3(\mathbb{O})$ since they can't be written as non-negative combinations of other idempotent matrices.

(c) Each face of $C_3(\mathbb{O})$ is generated by an idempotent matrix, because in any cone all the positive linear combinations of the same set of vectors generate the same face, so a diagonal matrix with non-negative entries generates the same face as a matrix with only zeros and ones.

(d) Any two idempotent matrices of trace 1 lie in a face generated by an idempotent matrix of trace 2, because they can be put simultaneously in the form

$$\begin{bmatrix} a & x & 0 \\ \overline{x} & b & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{ and these lie in the face generated by } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

(e) $A \in C_3(\mathbb{O})$ is an idempotent of trace 1 if and only if $I - A$ is an idempotent with trace 2. If A and B are idempotents of trace 1, then A lies in the face generated by $I - B$ if and only if B lies in the face generated by $I - A$. This duality and (d)

show that any two faces generated by idempotent matrices of trace 2 meet in a face generated by an idempotent matrix of trace 1.

2. FACE LATTICES DEFINING PROJECTIVE SPACES.

If C is a convex body whose face lattice is modular and irreducible and C is not strictly convex, the set of extreme points of C is a projective space with flats determined by the faces of C . We would like to know which projective spaces arise in this way, and what convex bodies give rise to them.

By Blaschke selection theorem [7], the space of all compact, convex subsets of a convex body in $\mathbb{R}^n$, with the Hausdorff metric, is compact. So the subspace formed by the compact convex subsets of the boundary is closed, but the subspace formed by the faces is not closed in general.

Lemma 2. *If C is a convex body whose face lattice $\mathcal{F}$ is modular, then the set $\mathcal{F}_h$ of faces of rank h , with the Hausdorff metric, is compact for each h .*

Proof. Let F_i be a sequence of faces of rank h . Then F_i has a subsequence F_{i_j} that is convergent in $\mathcal{C}$, and its limit is a compact convex set K contained in ∂C , so K generates a proper face F of C of some rank h' . We claim that $h' = h$ and $K = F$.

If the rank of F was less than h , there would be a face F^c of C of rank $n - h$ with $F^c \cap F = \phi$. As F and F^c are two disjoint compact sets in $\mathbb{R}^n$, there exists $\epsilon > 0$ such that the ϵ -neighborhoods of F and F^c in $\mathbb{R}^n$ are disjoint. But as $F_{i_j} \rightarrow K \subset F$ in the Hausdorff metric, then for sufficiently large j , F_{i_j} is contained in the ϵ -neighborhood of F , therefore $F_{i_j} \cap F^c = \phi$, but these 2 faces have ranks that add up to n , so they should meet, a contradiction.

If the rank of F is h and $K \neq F$, there is an extreme point $p \in F - K$, and there is a face F' of rank $n - h$ that meets F only at p , so $F' \cap K = \phi$ and the previous argument gives a contradiction.

To show that the rank of F cannot be larger than h , proceed inductively on $n - h$. As a limit of proper faces is contained in a proper face, the claim holds if $n - h = 1$. Given a sequence F_i of faces of rank h , let F be a face generated by the limit of a convergent subsequence F_{i_j} . If p is an extreme point of C not in F , then for sufficiently large j , $p \notin F_{i_j}$ (otherwise p would be in F). Let G_{i_j} be a face of rank $h + 1$ containing F_{i_j} and p . Now we can assume inductively that the limit of a convergent subsequence of G_{i_j} generates a face of rank $h + 1$. This face contains F properly (because it contains p) so $h' < h + 1$. $\square$

Now recall that a *topological projective space* is a projective space in which the sets of flats of each rank are given nontrivial topologies that make the join and meet operations $\vee$ and $\wedge$ continuous, when restricted to pairs of flats of fixed ranks whose join or meet have a fixed rank [5].

Lemma 3. *If C is a convex body whose face lattice is modular and irreducible then C is strictly convex or the set of extreme points $\mathcal{E}(C)$ is a topological projective space which is compact and connected.*

Proof. A natural topology for the set of flats is given by the Hausdorff distance between the faces. By lemma 2, $\mathcal{E} = \mathcal{F}_0$ is compact. As the lattice is irreducible and has more than 2 points, the 1-flats have more than 2 points, so (as they are topological spheres) they are connected. Now every pair of points in $\mathcal{E}$ is contained

in one of these spheres, so $\mathcal{E}$ is connected (one can actually show that each $\mathcal{F}_h$ is connected).

It remains to show that $\vee$ and $\wedge$ are continuous on the preimages of each $\mathcal{F}_h$. Suppose $A_i \rightarrow A$, $B_i \rightarrow B$ where $A_i \wedge B_i$ and $A \wedge B$ are faces corresponding to h flats. We need to show that $A_i \wedge B_i \rightarrow A \wedge B$. By lemma 2 $C_i = A_i \wedge B_i$ has convergent subsequences and the limit of a convergent subsequence C_{i_α} is a face C_α corresponding to an h flat. As C_{i_α} is contained in A_{i_α} and B_{i_α} , C_α is contained in $A \wedge B$. But C_α and $A \wedge B$ are both faces corresponding to h flats, so $C_\alpha = A \wedge B$. Similarly, if $A_i \rightarrow A$, $B_i \rightarrow B$ and $A_i \vee B_i$, $A \vee B$ are faces corresponding to h flats, the limit of each convergent subsequence of $D_i = A_i \vee B_i$ is a face D corresponding to an h flat. As D_i contains A_i and B_i , D contains $A \vee B$, and as both faces correspond to h flats they must be equal. $\square$

Let C be any convex body. Denote by $\mathcal{B}(C) \subset C$ the set of baricenters of faces of C and let $b : \mathcal{F}(C) \rightarrow \mathcal{B}(C)$ the function that assigns to each face its baricenter.

Lemma 4. (a) *If $\mathcal{F}(C)$ is compact then b is a homeomorphism.*

(b) *If $\mathcal{E}(C)$ is compact then a sequence of faces F_i converges to a face F if and only if $\mathcal{E}(F_i)$ converges to $\mathcal{E}(F)$.*

Proof. (a) The function that assigns to each compact convex set in $\mathbb{R}^n$ its baricenter is continuous, so $b : \mathcal{F}(C) \rightarrow \mathcal{B}(C)$ is a continuous bijective map from a compact Hausdorff space to a metric space.

(b) The Hausdorff distance between two compact convex sets is bounded above by the Hausdorff distance between their sets of extreme points.

If F_i converges to F but $\mathcal{E}(F_i)$ doesn't converge to $\mathcal{E}(F)$ then there is a subsequence $\mathcal{E}(F_{i_j})$ that stays at distance at least $\varepsilon > 0$ from $\mathcal{E}(F)$. For each i_j there is an extreme point $p_{i_j} \in F_{i_j}$ whose distance from $\mathcal{E}(F)$ is larger than ε , or an extreme point $q_i \in F$ whose distance from $\mathcal{E}(F_{i_j})$ is larger than ε . If there is a convergent subsequence $p_{i_k} \rightarrow p \in F$ then p is at distance at least ε from $\mathcal{E}(F)$, so p can't be an extreme point of C .

If there is a convergent subsequence $q_{i_k} \rightarrow q \in F$, take $p'_{i_k} \in F_{i_k}$ with $p'_{i_k} \rightarrow q$. Eventually $|p'_{i_k} - q_{i_k}| < \frac{\varepsilon}{2}$ so the distance from p'_{i_k} to $\mathcal{E}(F_{i_k})$ is at least $\frac{\varepsilon}{2}$, so p'_{i_k} is the center of a straight interval I_{i_k} of length ε contained in F_{i_k} . A convergent subsequence of these intervals yields a straight interval centered at q and contained in F , so q can't be an extreme point of C , contradicting the compactness of $\mathcal{E}(C)$. $\square$

Lemma 5. *If C and C' are convex bodies with $\mathcal{F}(C)$ and $\mathcal{F}(C')$ compact, then any continuous "face preserving" map $\varphi : \mathcal{E}(C) \rightarrow \mathcal{E}(C')$ extends naturally to a continuous map $\varphi : C \rightarrow C'$.*

Proof. φ determines a function $\Psi : \mathcal{F}(C) \rightarrow \mathcal{F}(C')$. Ψ is continuous because by lemma 4, $F_i \rightarrow F$ implies $\mathcal{E}(F_i) \rightarrow \mathcal{E}(F)$, then uniform continuity of φ on $\mathcal{E}(C)$ implies that $\varphi(\mathcal{E}(F_i)) \rightarrow \varphi(\mathcal{E}(F))$ so by definition $\mathcal{E}(\Psi(F_i)) \rightarrow \mathcal{E}(\Psi(F))$ and so $\Psi(F_i) \rightarrow \Psi(F)$. So φ can be extended to a continuous function $\varphi : \mathcal{B}(C) \rightarrow \mathcal{B}(C')$ as $b \circ \Psi \circ b^{-1}$ (recall that $\mathcal{E}(C) \subset \mathcal{B}(C)$). Now we can extend φ to the interiors of the faces of C defining it linearly on rays, as follows.

For each point $a \in C$, let $F(a)$ be the unique face of C containing a in its interior and let $b(a)$ be the baricenter of $F(a)$. Although $F(a)$ and $b(a)$ are not continuous functions of a on all of C , they are continuous on the union of the interiors of the faces corresponding to h -flats for each h . If $a \neq b(a)$ let $p(a)$ be the

projection of a to $\partial F(a)$ from $b(a)$ and let $\lambda(a) = \frac{|a-b(a)|}{|p(a)-b(a)|}$ (or 0 if $a = b(a)$) so $a = (1 - \lambda(a))b(a) + \lambda(a)p(a)$. Define $\varphi(a) = (1 - \lambda(a))\varphi(b(a)) + \lambda(a)\varphi(p(a))$.

Assume inductively that φ is continuous on the union of $\mathcal{B}(C)$ and the faces of C of dimension less than d (this set is closed because the limit of faces of dimension less than d has dimension less than d). and let's show that for each sequence of points a_i in the interiors of faces of dimension d , $a_i \rightarrow a$ implies $\varphi(a_i) \rightarrow \varphi(a)$. We may assume that the a_i are not baricenters, so $p(a_i)$ is well defined.

Case 1. $F(a_i) \rightarrow F(a)$ then $b(a_i) \rightarrow b(a)$ by the continuity of b on faces.

If $b(a) \neq a$ then $p(a_i) \rightarrow p(a)$ and $\lambda(a_i) \rightarrow \lambda(a)$ so $\varphi(a_i) = (1 - \lambda(a_i))\varphi(b(a_i)) + \lambda(a_i)\varphi(p(a_i)) \rightarrow (1 - \lambda(a))\varphi(b(a)) + \lambda(a)\varphi(p(a)) = \varphi(a)$.

If $b(a) = a$ then $\lim b(a_i) = \lim a_i$ but $p(a_i)$ may not converge, so consider a convergent subsequence $p(a_{i_j})$: If $\lim p(a_{i_j}) \neq \lim b(a_{i_j}) = \lim a_{i_j}$ then $\lim \lambda(a_{i_j}) = 0$ so $\varphi(a_{i_j}) = \varphi(b(a_{i_j})) + \lambda(a_{i_j})[\varphi(p(a_{i_j})) - \varphi(b(a_{i_j}))] \rightarrow \varphi(b(a)) + 0 = \varphi(a)$. If $\lim p(a_{i_j}) = \lim b(a_{i_j})$ then $\lim \varphi(p(a_{i_j})) = \lim \varphi(b(a_{i_j}))$ (by continuity of φ in the baricenters and faces of lower dimension) and as $\varphi(a_{i_j})$ lies between them, $\lim \varphi(a_{i_j}) = \lim \varphi(b(a_{i_j})) = \varphi(b(a)) = \varphi(a)$.

Case 2. $F(a_i) \nrightarrow F(a)$, then for any convergent subsequence $F(a_{i_j})$ with limit a face $F \neq F(a)$, a lies in F and so a must lie in ∂F , so $|a_{i_j} - p(a_{i_j})| \rightarrow 0$ and $\lambda(a_{i_j}) \rightarrow 1$, so $\lim \varphi(a_{i_j}) = \lim(1 - \lambda(a_{i_j}))\varphi(b(a_{i_j})) + \lambda(a_{i_j})\varphi(p(a_{i_j})) = \lim \varphi(p(a_{i_j})) = \varphi(a)$ (by continuity of φ on the faces of lower dimension). $\square$

Theorem 1. *Let C be a convex body whose face lattice defines a n -dimensional projective space.*

If $n = 2$, then C has dimension 5, 8, 14 or 26.

If $n > 2$ (or the space is desarguesian) there is a face-preserving homeomorphism from C to a section of a cone of Hermitian matrices over $\mathbb{R}$, $\mathbb{C}$, or $\mathbb{H}$.

Proof. First consider the case $n = 2$. All the lines of a topological projective plane $\mathcal{P}$ are homeomorphic because if l is a line and p is a point not in l then the projection $\phi : \mathcal{P} - p \rightarrow l$, $\phi(x) = (x \vee p) \wedge l$ is continuous and its restriction to each projective line not containing p is one to one. If the projective lines are topological spheres then a famous result of Adams [5, p.1278], shows that their dimension must be $d = 0, 1, 2, 4$ or 8 .

To compute the dimension of C take 3 faces of rank 1, F_0, F_1 and F_2 so that F_1 and F_2 meet at a point p not in F_0 . The projection $\phi : \mathcal{E}(C) - \{p\} \rightarrow \partial F_0$ extends to a continuous map $\phi : \cup \{F \mid F \text{ face of } C, p \notin F\} \rightarrow F_0$ whose restriction to each face is one to one (see proof of lemma 5). Now $U = \cup \{Int F \mid F \text{ face of } C, p \notin F\}$ is an open subset of ∂C and the function $\Phi : U \rightarrow F_0 \times (\partial F_1 - \{p\}) \times (\partial F_2 - \{p\})$ defined as $\Phi(x) = (\phi(x), \partial F(x) \wedge \partial F_1, \partial F(x) \wedge \partial F_2)$ is continuous and bijective, so U has the same dimension as $F \times \partial F \times \partial F$, which is $3d + 1$, therefore C has dimension $3d + 2$. Note that the discrepancy between the dimensions of the union of the boundaries of the faces ($2d$) and the union of the faces ($3d + 1$) arises because the boundaries of the faces overlap (as the lines in a projective plane do) but the interiors of the faces are disjoint. When $n > 2$, there is a similar homeomorphism from an open subset of ∂C and a product $F_0 \times (\partial F_1 - \{p\}) \times (\partial F_2 - \{p\}) \times \dots \times (\partial F_n - \{p\})$ where F_0 is a face of rank $r - 1$ and $F_1, F_2, \dots, F_n$ are faces of rank 1. So $\dim(C) = \dim(F_0) + rd + 1$ and it follows by induction that $\dim C = \frac{n(n-1)}{2}d + n - 1$.

Now assume that the projective space determined by $\mathcal{E}(C)$ is desarguesian. Every topological desarguesian projective space is isomorphic to a projective space over a

topological division ring A (defined on a line minus a point) and the isomorphism is a homeomorphism [5, p.1261]. By a classic result of Pontragin [5, p.1263] the only locally compact, connected division rings are $\mathbb{R}$, $\mathbb{C}$ and $\mathbb{H}$. So the projective space determined by $\mathcal{E}(C)$ is isomorphic to $\mathbb{RP}^r$, $\mathbb{CP}^r$ or $\mathbb{HP}^r$. Therefore $\mathcal{E}(C)$ is isomorphic to $\mathcal{E}(C')$ where C' is a section of a cone of Hermitian matrices, and so by 5 there is a face-preserving homeomorphism from C to C' . $\square$

3. FACE LATTICES DEFINING AFFINE SPACES.

Now let us consider closed (but not necessarily compact) convex sets in $\mathbb{R}^n$ whose faces meet as the subspaces of an affine space. An abstract affine plane consists of a set of points and a set of lines so that 1) there are at least 2 points and 2 lines 2) every pair of points is contained in a line and 3) given a line and a point not contained in it, there is a unique line containing the point and parallel to the line. The axioms of an abstract affine space are not so simple, but it is enough to know that if P is a projective space then the complement of a maximal flat of P is an affine space, and any affine space A can be embedded in a projective space in this fashion, by attaching to A a point at infinity for each parallelism class of affine lines.

Observe that if a closed convex set C in $\mathbb{R}^n$ is non-compact, it contains a ray (half of a euclidean line) and if C contains a ray then it contains all the parallel rays starting at points of C (we say that C contains an infinite direction). So if C contains a line, C is the product of that line and a closed convex set C' of lower dimension and the face lattice of C and C' are isomorphic. So from now on we will assume that C doesn't contain lines.

It is easy to see that the faces of a polytope cannot determine an affine space (the faces of rank i would have dimension i , two parallel faces of rank 1 generate a face of rank 2 with at least 4 vertices, but the sides of a polygon don't define an affine plane).

Example 3. *Let C be a convex body in $\mathbb{R}^n$ whose faces determine a projective space. Take a cone over C and slice it with a hyperplane parallel to a support hyperplane containing a maximal face. The result is a closed non-compact convex set C' in $\mathbb{R}^n$ whose faces determine an affine space. In particular, the cones of Hermitian matrices have non-compact sections whose face lattice determines a real, complex or quaternionic affine space or an octonionic affine plane.*

$\mathbb{RP}^n$ can be seen as the space of lines through the origin in $\mathbb{R}^{n+1}$ or as the quotient of the unit sphere $\mathbb{S}^n$ (or the sphere at infinity of $\mathbb{R}^{n+1}$) by the action of the antipodal map. Identifying $\mathbb{R}^n$ with a hyperplane of $\mathbb{R}^{n+1}$ that doesn't contain the origin gives an embedding of $\mathbb{R}^n$ as a dense open subset of $\mathbb{RP}^n$. The remaining points of $\mathbb{RP}^n$ correspond to lines through the origin in $\mathbb{R}^{n+1}$ that don't meet the hyperplane, i.e., parallelism classes of lines in $\mathbb{R}^n$ (or pairs of antipodal points in the sphere at infinity). Define a set in $\mathbb{RP}^n$ to be *convex* if it is the image of a convex set in $\mathbb{R}^n$ under one of these embeddings. As convex sets in $\mathbb{RP}^n$ correspond to convex cones based at the origin of $\mathbb{R}^{n+1}$, convexity in $\mathbb{RP}^n$ doesn't depend on the particular embedding, and a convex set in $\mathbb{RP}^n$ has the usual properties of a convex set in $\mathbb{R}^n$.

Now if C is a closed convex set in $\mathbb{R}^n$ that doesn't contain lines, its closure $\overline{C}$ is a convex set in $\mathbb{RP}^n$. The faces of $\overline{C}$ are the closures of faces of C and their intersections with the sphere at infinity modulo the antipodal map.

Lemma 6. *In a closed convex set in $\mathbb{R}^n$ that has semi-modular face lattice, two faces of rank 1 can share at most one direction, and it corresponds to a ray.*

Proof. We are considering convex bodies without lines. Suppose that two rank 1 faces F_1 and F_2 have a common direction, i.e., there are segments of parallel euclidean lines l_1 and l_2 lying in F_1 and F_2 . We may assume that l_i goes through an interior point of F_i , so l_i meets ∂F_i in one or two extreme points. If l_1 or l_2 has two extreme points then there is a convex quadrilateral with sides in l_1 and l_2 with 3 extreme points as vertices. The interior of the quadrilateral lies in the interior of the rank 2 face generated by the 3 extreme points, but the intersection of its diagonals lies in the rank 1 face generated by 2 extreme points, a contradiction. So l_i meets ∂F_i in only one point and so F_i contains a ray l_i^+ . If F_1 and F_2 have two common directions, there is a common direction which meets F_1 in 2 points, giving the same contradiction. $\square$

Lemma 7. *If the faces of a closed convex set C in $\mathbb{R}^n$ define an affine space, then each face representing a line contains a unique ray, and faces representing parallel lines contain parallel rays.*

Proof. We are assuming again that C doesn't contain lines, so the points of the affine space correspond to the extreme points of C . Each affine lines is represented by the boundary of a convex set of dimension at least 2, which is connected, so set of extreme points in C is connected.

Let's first show that the faces representing affine lines cannot be compact. Suppose that C has a compact face F . Let p and q be two extreme points in F and let q_i be a sequence of extreme points not in F that converge to q . Let F_i be the face generated by p and q_i . If F_i is non-compact, it contains a ray r_i through p . So F_i contains the "parallelogram" determined by the interval pq_i and the ray r_i . An infinite sequence of l_i 's would have a subsequence converging to a ray l through q , so the parallelogram determined by the interval pq and the ray l would be contained in ∂C , so it would have to be contained in a face of C , which would have to be F because it contains p and q . This contradicts the assumption that F is compact and shows that if q_i is sufficiently close to q , the face generated by p and q_i is compact. Now take a face F' that doesn't meet F (i.e., F and F' represent parallel affine lines). As F is compact and F' is closed in $\mathbb{R}^n$, there is an ε neighborhood of F that doesn't intersect F' . By the previous argument there is a point q_i not in F so that the face F_i generated by p and q_i is contained in the ε neighborhood of F . So F_i doesn't meet F' , but F was supposed to be the only face containing p and disjoint from F' .

This proves that F is non-compact, so it contains rays. Let's show that two faces representing parallel affine lines contain parallel rays. Let p be an extreme point outside F , so F and p generate a face H representing an affine plane. There are extreme points $p_0, p_1, p_2, \dots$ in F so that the sequence of intervals p_0p_i converges to a ray l_+ contained in F (because F is closed). The sequence of intervals pp_i lie in ∂H and converge to a ray m_+ parallel to l_+ and containing p so (as H is closed) m_+ is contained in a face G of ∂H representing an affine line. As two faces that contain parallel rays cannot meet at a single point, G doesn't meet F so (as F and G are contained in H) G represents the affine line parallel to F through p . If F has nonparallel rays, one can construct as before two nonparallel rays l_+ and l'_+ in F and faces G and G' through p and containing rays m_+ and m'_+ . The uniqueness of

parallel affine lines implies that $G = G'$, so F and G have more than one common direction, contradicting the previous lemma. $\square$

This shows that if the faces of a closed convex set C define an affine space, C is non-compact. One can show that if the faces of a closed convex set C (containing no lines) define a projective space, C must be compact. For this, one has to give a topology to the space of closed convex sets in $\mathbb{R}^n$ that makes it locally compact, show that this makes $\mathcal{E}(C)$ into a locally compact projective space, and observe that these spaces are necessarily compact.

Theorem 2. *If C is a closed convex set in $\mathbb{R}^n$ whose faces determine an affine space, there is a projective transformation in $\mathbb{RP}^n$ taking $\overline{C}$ to a compact convex set in $\mathbb{R}^n$ whose faces determine a projective space. If the space is desarguesian, there is a face-preserving homeomorphism from C to a non-compact slice of a cone of Hermitian matrices.*

Proof. We need to show that if the face lattice of C determines an affine space, the face lattice of $\overline{C} \subset \mathbb{RP}^n$ determines its projective completion. The faces of C representing affine lines are non-compact, and two of them share an infinite direction in $\mathbb{R}^n$ if and only if they represent parallel affine lines. The closure $\overline{C} \subset \mathbb{RP}^n$ contains one point at infinity for each infinite direction in C , so $\overline{C}$ contains an extreme point at infinity for each class of faces of C representing parallel lines. This corresponds precisely with the definition of the projective completion of the affine space. Now the result for C follows by applying theorem 1 to $\overline{C}$. $\square$

4. PROJECTIVE PLANES AND THE CASE $d = 1$.

The face lattice of a convex body C (not a triangle) determines a projective plane if every pair of extreme points is contained in a proper face and every pair of faces with more than one point meet. By theorem 1 this projective plane is compact and connected, so for some $d \in \{1, 2, 4, 8\}$, all the faces of C have dimension $d + 1$ and C has dimension $n = 3d + 2$.

Lemma 8. *A $d + 1$ dimensional subspace of $\mathbb{R}^n$ is the span of a face of C if and only if it meets all the spans of faces of C .*

Proof. Let S be an affine subspace that intersects $\text{span}(F)$ for every $F \in \mathcal{F}_1$, the set of faces of rank 1. Then $\{F \in \mathcal{F}_1 \mid \dim(S \cap \text{span}(F)) \geq i\}$ is closed in $\mathcal{F}_1$ for each i .

Case 1. $d = 1$. We claim that if S is not the span of a face then S cannot intersect $\text{span}(F)$ in more than one point. For, if $S \cap \text{span}(F)$ contains a line, then $\text{span}(S \cup F)$ is 3 dimensional. Take an extreme point $p \notin \text{span}(S \cup F)$ and let F_1 and F_2 be 2 faces containing p , and meeting F at points p_1 and p_2 not in S . If p'_1 and p'_2 are points in $S \cap \text{span}(F_1)$ and $S \cap \text{span}(F_2)$ respectively, then p, p_i and p'_i are not aligned (otherwise p would be in the span of $S \cup \text{span}(F)$) and so the span of p, p_i and p'_i , which is $\text{span}(F_i)$, is contained in $\text{span}(p \cup S \cup F)$. But $\text{span}(p \cup S \cup F)$ is 4 dimensional, so it cannot contain the 3 faces F, F_1 and F_2 because if it did, it would contain each face that meets F, F_1 and F_2 at 3 different points, but every face is a limit of such faces, so it would contain all the faces of C , but C has dimension 5. This shows that S intersects each $\text{span}(F)$ at exactly one point, and so S contains at most one extreme point of C . The function $I : \mathcal{F}_1 \rightarrow S$ that maps each face F_i to the point of intersection of $\text{span}(F)$ with S is continuous, and

as the spans of faces meet only at extreme points, I only fails to be injective on the faces containing the extreme point in S (if any). But $\mathcal{F}_1$ is a 2-dimensional closed surface in which the faces that contain an extreme point form a closed curve, and there are no continuous maps from a closed surface to the plane that fail to be injective only along a curve.

Case 2. S doesn't contain extreme points of some face F . Choose F that minimizes the dimension of the subspace $S \cap \text{span}(F)$. Then for every F' in a neighborhood of F , $S \cap \text{span}(F')$ is a subspace of minimal dimension and with no extreme points. If S' is the orthogonal complement of $S \cap \text{span}(F)$ in S then S' intersects $\text{span}(F)$ in one point for all F' in a smaller neighborhood V of F . Then the function $I : V \rightarrow S'$ that maps F' to $S' \cap \text{span}(F')$ is continuous, and it is injective as the spans of faces only meet at extreme points. But an injective map between manifolds can only exist when the domain has dimension no larger than the target so $2d \leq \dim S' \leq \dim S \leq d + 1$, so $d = 1$ and we are in case 1.

Case 3. S contains extreme points of each face F . As C is convex, either $S \cap C \subset \partial C$ or $\partial_S(S \cap C) = S \cap \partial C$. In the first case $S \cap C$ is contained in a face F_1 of C and so either $S \cap C = F_1$ (so $F_1 \subset S$) or there is an extreme point p of F_1 not contained in S , but then a face F_2 that meets F_1 at p doesn't meet $F_1 \cap S \supset S \cap C$ so S doesn't contain extreme points of F_2 .

Let p be an extreme point not in S , and consider the set $\mathcal{F}_1^p$ of faces of rank 1 containing p . If F and F' are distinct faces in $\mathcal{F}_1^p$, $S \cap F$ and $S \cap F'$ are disjoint. Choose F so that $S \cap F$ has minimal dimension, then for all F' in some neighborhood V of F , $S \cap F'$ has the same dimension and the map $I_B : \mathcal{F}_1^p \cap V \rightarrow S \cap \partial C$ that sends F' to the baricenter of $S \cap F'$ is continuous and injective. As I_B is a map between manifolds, $d = \dim \mathcal{F}_1^p \leq \dim S \cap \partial C \leq d$ and so by domain invariance the image of I_B is an open subset of $S \cap \partial C = \partial_S(S \cap C)$. This implies that for each $F' \in \mathcal{F}_1^p \cap V$, $S \cap F'$ consists of one point (if a face of a convex set has more than 1 point, its baricenter is arbitrarily close to points in the boundary that are not baricenters of other faces, namely, the points in the face) and so, by hypothesis, $S \cap F'$ is an extreme point of C .

So part of the boundary of $S \cap C$ in S is strictly convex, therefore the line segment joining two extreme points in it lies in $\text{Int}_S(S \cap C)$, but that line segment lies in the face of C containing the 2 extreme points, so it must lie in $S \cap \partial C = \partial_S(S \cap C)$, a contradiction. $\square$

Lemma 9. *The boundaries of the faces of rank 1 of C are semi-algebraic sets. If $d = 1$, they are conic sections.*

Proof. By lemma 8, the set $\mathcal{S}$ of spans of faces of C is the same as the set of $d + 1$ -dimensional subspaces of $\mathbb{R}^n$ that intersect every element of $\mathcal{S}$. The set of all $d + 1$ -dimensional affine subspaces of $\mathbb{R}^n$ forms a real algebraic variety and the condition that the subspaces meet a fixed subspace is algebraic, so (by the finite descending chain condition) there is a finite family of spans $S_1, S_2, \dots, S_m \in \mathcal{S}$ such that $S \in \mathcal{S}$ if it intersects these S_i 's (see [1]).

Now for $(x_1, x_2, \dots, x_m) \in S_1 \times S_2 \dots \times S_m$, the subspace $\text{span}(x_1, \dots, x_m)$ has dimension at least $d + 1$ (otherwise it would be contained in two subspaces of dimension $d + 1$ that meet each S_i , so they would both be in $\mathcal{S}$, but two spans can only meet in 1 point). So $\text{span}(x_1, \dots, x_m)$ lies in $\mathcal{S}$ if and only if its dimension is $d + 1$, and this happens if and only if some determinants (given by polynomials on $x_1, \dots, x_m$) vanish. Therefore the set $X = \{(x_1, x_2, \dots, x_m) \in S_1 \times S_2 \dots \times S_m \mid \text{span}(x_1, \dots, x_m) \in \mathcal{S}\}$ is

real algebraic, as is the set X^p formed by the elements of X that contain a fixed point p . If F_1 is the face in S_1 and p is an extreme point of C outside F_1 then ∂F_1 consists of the intersections of S_1 with the elements of $\mathcal{S}$ containing p . So ∂F_1 is the one to one projection of the algebraic set X^p to S_1 , so ∂F_1 is at least semi-algebraic.




Figure 1

Now assume $d = 1$ so $n = 5$. Every projective plane has 7 points and 6 lines so that each line contains 3 points as in figure 1, so C has 7 extreme points and 6 faces intersecting in that way. The 7 points are in general position in $\mathbb{R}^5$ because as each face of C is spanned by 3 points, the span of any 6 of those points contains the span of 3 faces, which is all of $\mathbb{R}^5$. Therefore we may assume (by applying a projective transformation) that the 7 points are $p_0 = (0, 0, 0, 0, 0)$, $p_1 = (1, 0, 0, 0, 0)$, ..., $p_5 = (0, 0, 0, 0, 1)$, $p_6 = (1, 1, 1, 1, 1)$. Let S_i be the plane spanned by the face F_i . A plane S that intersects S_1, S_2 and S_3 has a parametrization $(x, y, z, v, w) = r(a, b, 0, 0, 0) + s(0, 0, c, d, 0) + t(e, e, e, e, f)$ with $r + s + t = 1$. S intersects S_4, S_5 and S_6 only if three systems of linear equations in r, s, t represented by the following matrices have nontrivial solutions:

$$\begin{vmatrix} a & 0 & e \\ 0 & d & e \\ b-1 & c-1 & 2e+f-1 \end{vmatrix} \begin{vmatrix} b & 0 & e \\ 0 & c & e \\ a-1 & d-1 & 2e+f-1 \end{vmatrix} \begin{vmatrix} b & 0 & e-f \\ 0 & d & e-f \\ a-1 & c-1 & 2e-f-1 \end{vmatrix}$$

As the determinants of these matrices are linear functions on the variables e and f , they vanish simultaneously if and only if the matrix of this new system has determinant 0:

$$\det \begin{vmatrix} -ac + 2ad - bd + a + d & ad & -ad \\ -ac + 2bc - bd + b + c & bc & -bc \\ -ad - bc + 2bd + b + d & ad + bc - bd - b - d & -bd \end{vmatrix} = 0$$

This determinant factors as the product of a linear and a quadratic function of a and b (with coefficients in c and d). Since the boundary of the face F_1 is formed by the intersections of S_1 with the planes that meet all S_i 's and go through a fixed point in the boundary of F_2 (this corresponds to fixing c and d), the boundary of F_1 is contained in the union of a line and a conic. As the boundary of F_1 is strictly convex, it must be the conic. $\square$

Theorem 3. *All convex bodies in $\mathbb{R}^5$ with modular and irreducible face lattice are projectively equivalent.*

Proof. Let C and C' be two such bodies. Take extreme points $p_0, p_1, \dots, p_6$ and faces $F_1, \dots, F_6$ of C as in figure 1. Pick an extreme point p'_0 in C' and two faces F'_1 and F'_2 of C' intersecting at p'_0 . Let S_i be the span of F_i . As the faces of C and C' are conics, there are linear transformations from S_1 to S'_1 taking F_1 to F'_1 and from S_2 to S'_2 taking F_2 to F'_2 . Together, they define a linear transformation l from $\text{span}(F_1 \cup F_2)$ to $\text{span}(F'_1 \cup F'_2)$. Let $p'_i = l(p_i)$ for $i = 1, \dots, 4$. The faces F_4, F_5, F_6 are generated by unique pairs of p_i 's with $i \leq 4$. Let F'_4, F'_5, F'_6 be the faces generated by the corresponding pairs of p'_i 's. Finally, let $p'_5 = S'_4 \cap S'_5$, let F'_3 be the face generated by p'_0 and p'_5 and let $p'_6 = S'_3 \cap S'_6$. The linear transformation l can be extended to a projective transformation ρ in $\mathbb{RP}^5$ that takes p_5 to p'_5 and p_6 to p'_6 . As ρ sends each p_i to p'_i , it sends each S_i to S'_i , so it sends each plane in $\mathbb{R}^5$ intersecting every S_i to a plane intersecting every S'_i . Since by construction ρ takes those planes that meet ∂F_1 and ∂F_2 to planes that meet $\partial F'_1$ and $\partial F'_2$, lemma 8 implies that ρ maps spans of faces of C to spans of faces of C' and therefore it maps faces to faces. $\square$

Question 1: *Are all the convex bodies whose face lattices determine classical projective spaces projectively equivalent to sections of cones of hermitian matrices?*

Question 2: *Can two convex bodies of the same dimension define non isomorphic projective planes (so they are not related by a face-preserving homeomorphism)?*

In dimensions 8 and 14 this is equivalent to ask if the projective planes are always desarguesian. In dimension 26 there might be enough space for non-equivalent non-desarguesian examples.

REFERENCES

- [1] Bochnak, J., Coste, M., Roy, M., Real Algebraic geometry. *Springer*, 1998.
- [2] Baez, J., The Octonions, *Bull. Amer. Math. Soc.* **39** (2001), 145-205.
- [3] Barker, G.P., Theory of cones. *Linear Algebra Appl* **39** (1981), 263-291.
- [4] Birkhoff, G., Lattice Theory, 3rd edition, *Amer. Math. Soc. Colloq. Publ.*, AMS 1967.
- [5] Grundhofer, T., Lowen, R., Linear topological geometries, in Handbook of Incidence Geometry. Buildings and Foundations. *North-Holland*. 1995.
- [6] Crawley, P., Dilworth R.P., Algebraic Theory of Lattices, *Prentice-Hall, Inc.*, 1973.
- [7] Gruber, P.M., Willis, J.M., Handbook of Convex Geometry. Volume A. *North-Holland*. 1993.
- [8] Hartshorne, R., Foundations of Projective Geometry. *W.A. Benjamin, Inc.* 1967.
- [9] Labardini-Fragoso, D., The face lattice of a finite dimensional cone, *Bachelor Thesis UNAM*, 2004.
- [10] Salzmann, H., Betten, D., Grundhofer, T., Haehl, H., Loewen, R., Stroppel, M., Compact Projective Planes. *De Gruyter Expositions in Mathematics*. 1995.

GRADUATE PROGRAM IN MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA.

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CUERNAVACA, MÉXICO.